



eCAMPUS
LEARN

CAMPUS HANDBOOK

IVO DE PAUW BIEKE MASSELIS

Animation Maths

Lannoo
Campus

eCampusLearn



eCAMPUS
LEARN

Go to www.ecampuslearn.com.

Fill in the following code:

Good luck!

Chapter 1, David Ritter; 2, John Evans; 3, Wouter Verweirder; 4, Daryl Beggs, Juan Pablo Arancibia Medina; 5, Stephanie Berghaeuser; 6, Martin Walls; 7, 12, Wouter Tansens; 8, Danie Pratt; 9, Ivo De Pauw; 10, Caetano Lacerda; 11, Ken Munyard; 13, Bieke Masselis; 14, Boke Haide; 15, Waldemar Zielinski; 16, Detje Holger; 17, Cornelia Roessing; p.25, p.103, Wouter Tansens; p.48, Wouter Verweirder; p.52, Leo Storme; p.185, Bieke Masselis; p.268, Yu-Sung Chang; p.343, Angelo Fallein.

D/2021/45/69 – ISBN 978 94 014 7495 5 – NUR 918

Cover design: Keppie & Keppie

Interior design: Ivo De Pauw, Bieke Masselis

© Ivo De Pauw, Bieke Masselis & Lannoo Publishers nv, Tiel, 2021.

LannooCampus Publishers is a subsidiary of Lannoo Publishers,
the book and multimedia division of Lannoo Publishers nv.

All rights reserved.

No part of this publication may be reproduced and/or made public,
by means of printing, photocopying, microfilm or any other means,
without the prior written permission of the publisher.

LannooCampus Publishers

Vaartkom 41 box 01.02

3000 Leuven

Belgium

www.lannoocampus.com

P.O. Box 23202

1100 DS Amsterdam

Netherlands

This book is dedicated to Malaika.

*“Sometimes I’m black, sometimes I’m white
it all depends on who is on the other side
there are things they cannot see
and there are things I can not hide”*

Bruno Deneckere (Someday, June 2006)

Contents

<i>Acknowledgements</i>	15
Chapter 1 · Arithmetic refresher	<u>17</u>
1.1 <i>Algebra</i>	18
Real numbers	18
Real polynomials	23
1.2 <i>Equations in one variable</i>	25
Linear equations	25
Quadratic equations	26
1.3 <i>Logarithms</i>	31
1.4 <i>Exercises</i>	34
Chapter 2 · Linear systems	<u>37</u>
2.1 <i>Definitions</i>	38
2.2 <i>Methods for solving linear systems</i>	40
Solving by substitution	40
Solving by elimination	41
2.3 <i>Exercises</i>	45
Chapter 3 · Trigonometry	<u>47</u>
3.1 <i>Angles</i>	48
3.2 <i>Triangles</i>	50
3.3 <i>Right triangle</i>	54
3.4 <i>Unit circle</i>	55
3.5 <i>Special angles</i>	57
Trigonometric ratios for an angle of $45^\circ = \frac{\pi}{4}$ rad	58
Trigonometric ratios for an angle of $30^\circ = \frac{\pi}{6}$ rad	58
Trigonometric ratios for an angle of $60^\circ = \frac{\pi}{3}$ rad	59
Overview	59
3.6 <i>Pairs of angles</i>	60
3.7 <i>Sum identities</i>	60
3.8 <i>Inverse trigonometric functions</i>	63
3.9 <i>Exercises</i>	65

Chapter 4 · Functions	<u>67</u>
4.1 <i>Basic concepts on real functions</i>	68
4.2 <i>Polynomial functions</i>	69
Linear functions	69
Quadratic functions	71
4.3 <i>Intersection of functions</i>	73
4.4 <i>Logarithmic functions</i>	75
4.5 <i>Exponential functions</i>	76
4.6 <i>Trigonometric functions</i>	78
Elementary sine function	78
General sine function	78
Transversal oscillations	82
4.7 <i>Inverse trigonometric functions</i>	82
arcsine	82
arccosine	84
arctangent	84
atan2	85
4.8 <i>Maclaurin expansions</i>	86
4.9 <i>Exercises</i>	89
Chapter 5 · The Golden Section	<u>91</u>
5.1 <i>The golden number</i>	92
5.2 <i>The golden section</i>	94
The golden triangle	94
The golden rectangle	95
The golden spiral	96
The golden pentagon	98
The golden ellipse	98
5.3 <i>Golden arithmetic</i>	99
Golden identities	99
The Fibonacci numbers	100
5.4 <i>The golden section worldwide</i>	103
5.5 <i>Exercises</i>	105
Chapter 6 · Coordinate systems	<u>107</u>
6.1 <i>Cartesian coordinates</i>	108
6.2 <i>Parametric curves</i>	108
6.3 <i>Polar coordinates</i>	112
6.4 <i>Polar curves</i>	115
6.5 <i>Exercises</i>	118

Chapter 7 · Vectors	<u>121</u>
7.1 <i>The concept of a vector</i>	122
Vectors as arrows	122
Vectors as arrays	123
Free vectors	126
Base vectors	126
7.2 <i>Addition of vectors</i>	127
Vectors as arrows	127
Vectors as arrays	127
Vector addition summarised	128
7.3 <i>Scalar multiplication of vectors</i>	129
Vectors as arrows	129
Vectors as arrays	129
Scalar multiplication summarised	130
Normalisation	130
Properties	130
7.4 <i>Vector subtraction</i>	131
Creating free vectors	131
Euler's method for trajectories	132
7.5 <i>Decomposition of vectors</i>	133
Decomposition of a plane vector	133
Base vectors defined	134
7.6 <i>Dot product</i>	135
Definition	135
Geometric interpretation	136
Orthogonality	139
7.7 <i>Cross product</i>	140
Definition	140
Geometric interpretation	142
Parallelism	144
7.8 <i>Normal vectors</i>	145
7.9 <i>Exercises</i>	147

Chapter 8 · Parameters	<u>149</u>
8.1 Parametric equations	150
8.2 Vector equation of a line	151
8.3 Intersecting straight lines	155
8.4 Vector equation of a plane	157
8.5 Exercises	161
Chapter 9 · Kinematics	<u>163</u>
9.1 Measures	164
9.2 Deltatime	165
9.3 Translational motion	165
Rectilinear motion with constant velocity (RMCV)	168
Rectilinear motion with constant acceleration (RMCA)	168
Free fall	171
Summary	174
9.4 Circular motion	176
Uniform circular motion (UCM)	176
Nonuniform circular motion (NCM)	183
Summary	186
9.5 Planar curvilinear motion	187
Normal-tangential components	188
Radial-angular components	191
9.6 Independence of Motion	194
Combined rectilinear motions with constant velocities	194
Projectile motion (PM)	195
9.7 Exercises	200
Chapter 10 · Collision detection	<u>203</u>
10.1 Collision detection using circles and spheres	204
Circles and spheres	204
Intersecting line and circle	206
Intersecting circles and spheres	208
10.2 Collision detection using vectors	211
Location of a point with respect to other points	211
Altitude to a straight line	212
Altitude to a plane	214
Frame rate issues	216
Location of a point with respect to a polygon	217
10.3 Exercises	220

Chapter 11 · Matrices	<u>223</u>
11.1 <i>The concept of a matrix</i>	224
11.2 <i>Determinant of a square matrix</i>	225
11.3 <i>Addition and scalar multiplication of matrices</i>	227
Addition of matrices	227
Scalar multiplication of a matrix	229
11.4 <i>Transpose of a matrix</i>	230
11.5 <i>Dot product of matrices</i>	230
Introduction	230
Condition	232
Definition	232
Properties	233
11.6 <i>Inverse of a matrix</i>	235
Introduction	235
Definition	235
Conditions	236
Row reduction	236
Matrix inversion	237
Inverse of a product	240
Solving systems of linear equations	241
11.7 <i>The Fibonacci operator</i>	243
11.8 <i>The matrix exponential</i>	245
Structures	245
Matrix exponential	245
11.9 <i>Exercises</i>	247
 Chapter 12 · Bezier curves	 <u>249</u>
12.1 <i>Vector equation of segments</i>	250
Linear Bezier segment	250
Quadratic Bezier segment	251
Cubic Bezier segment	252
Bezier segments of higher degree	254
12.2 <i>De Casteljau algorithm</i>	255
12.3 <i>Bezier curves</i>	256
Concatenation	256
Linear transformations	258
Illustrations	258
12.4 <i>Matrix representation</i>	260
Linear Bezier segment	260
Quadratic Bezier segment	261

Cubic Bezier segment	262
12.5 <i>B-splines</i>	264
Cubic B-splines	264
Matrix representation	265
De Boor's algorithm	267
12.6 <i>Exercises</i>	269
Chapter 13 · Transformations	<u>271</u>
13.1 <i>Translation</i>	272
13.2 <i>Scaling</i>	277
13.3 <i>Rotation</i>	280
Rotation in 2D	280
Rotation in 3D	282
13.4 <i>Reflection</i>	284
13.5 <i>Shearing</i>	286
13.6 <i>Combining standard transformations</i>	288
2D rotation around an arbitrary centre	290
3D scaling about an arbitrary centre	293
2D reflection over an axis through the origin	294
2D reflection over an arbitrary axis	295
3D combined rotation	298
13.7 <i>Row-representation</i>	299
13.8 <i>Exercises</i>	300
Chapter 14 · Transformation Analysis	<u>303</u>
Typesetting	304
14.1 <i>Translation analysis</i>	304
14.2 <i>Scaling analysis</i>	308
14.3 <i>Rotation analysis</i>	311
14.4 <i>Composite transformation analysis</i>	315
14.5 <i>Conventions</i>	318
14.6 <i>Applications</i>	319
Pivot transformation	319
Orbit transformation	320
Look-at transformation	322
14.7 <i>Exercises</i>	326
Chapter 15 · Scene Graphs	<u>329</u>
15.1 <i>Concept of a scene graph</i>	330
15.2 <i>Bone structures</i>	332

15.3	<i>Solar systems</i>	336
15.4	<i>Exercises</i>	339
Chapter 16	· View Transformation	<u>341</u>
16.1	<i>The Rendering Pipeline</i>	342
	The concept of the pipeline	342
	The stages of the pipeline	342
16.2	<i>Camera transformation</i>	344
16.3	<i>View transformation</i>	346
16.4	<i>View operator</i>	349
16.5	<i>Camera with zoom</i>	351
16.6	<i>Exercises</i>	354
Chapter 17	· Hypercomplex numbers	<u>357</u>
17.1	<i>Complex numbers</i>	358
17.2	<i>Complex number arithmetic</i>	362
	Complex conjugate	362
	Addition and subtraction	362
	Multiplication	363
	Exponentiation	365
	Division	366
17.3	<i>Complex numbers and transformations</i>	368
	Translation	368
	Standard rotation	369
	Standard scaling	370
	Composite transformation	370
17.4	<i>Complex continuation of the Fibonacci numbers</i>	371
	Integer Fibonacci numbers	371
	Complex Fibonacci numbers	372
17.5	<i>Quaternions</i>	374
17.6	<i>Quaternion arithmetic</i>	375
	Addition and subtraction	375
	Scalar multiplication	376
	Normalisation	376
	Quaternion multiplication	376
	Quaternion conjugate	378
	Inverse quaternion	379
17.7	<i>Quaternions and rotations</i>	379
	Trigonometrical representation of quaternions	380
	Euler representation of quaternions	380

Quaternion exponentiation	381
The quest for the quaternion rotation	382
Unit rotation quaternion	385
17.8 Exercises	389
Annex A · Real numbers in computers	<u>391</u>
A.1 Scientific notation	391
A.2 The decimal computer	391
A.3 Special values	392
Annex B · Notations and Conventions	<u>393</u>
B.1 Alphabets	393
Latin alphabet	393
Greek alphabet	394
B.2 Mathematical symbols	394
Sets	394
Mathematical symbols	395
Mathematical keywords	396
Numbers	396
Annex C · The International System of Units (SI)	<u>397</u>
C.1 SI Prefixes	397
C.2 SI Base measures	398
C.3 SI Supplementary measure	398
C.4 SI Derived measures	399
Bibliography	400
Index	403

Acknowledgements

We hereby insist on thanking a lot of people who made this book possible: Prof. Dr. Leo Storme, Wim Serras, Wouter Tansens, Wouter Verweirder, **Koen Samyn** [14] (credited for the chapters Transformation Analysis, Scene Graphs, View Transformation), Hilde De Maesschalck, Ellen Deketele, Conny Meuris, Hans Ameel, Dr. Rolf Mertig, Dick Verkerk, ir. Gose Fischer, Prof. Dr. Fred Simons, Sofie Eeckeman, Dr. Luc Gheysens, Dr. Bavo Langerock, Wauter Leenknecht, Marijn Verspecht, Sarah Rommens, Prof. Dr. Marcus Greferath, Dr. Cornelia Roessing, Tim De Langhe, Niels Janssens, Peter Flynn, Jurgen Leemans, Stef Lantsoght, Hilde Vanmechelen, Jef De Langhe, Ann Deraedt, Rita Vanmeirhaeghe, Prof. Dr. Jan Van Geel, Dr. Ann Dumoulin, Bart Uyttenhove, Rik Leenknecht, Peter Verswyvelen, Roel Vandommele, ir. Lode De Geyter, Bart Leenknecht, Olivier Rysman, ir. Johan Gielis, Frederik Jacques, Kristel Balcaen, ir. Wouter Gevaert, Bart Gardin, Dieter Roobrouck, Dr. Yu-Sung Chang (*Wolfram Demonstrations* [24]), Prof. Dr. Sy Blinder (*Wolfram Demonstrations*), Steven De Keninck, Prof. Dr. Mark McClure (*Wolfram Demonstrations*), Dr. Felipe Dimer de Oliveira (*Wolfram Demonstrations*), Steven Verborgh, Ingrid Viaene, Kayla Chauveau, Angelika Kirkorova, Thomas Vanhoutte, Fries Carton, Jef Daels, Andries Geens, Angelo Fallein, Jolan Plaum, Charles Derre, Anna Rich and whomever we might have forgotten!

Hereby our special thanks to **Dick Verkerk** for supporting our *Wolfram Application Server* [25]) server in the Netherlands at `can.nl`

The logo for 'Can' is displayed in a large, bold, black sans-serif font. The letter 'C' is significantly larger than the 'a' and 'n', and the 'a' has a unique shape with a small loop at the top.

Chapter 1 · Arithmetic refresher



As this chapter offers all the necessary mathematical skills for the full mastery of all further topics explained in this book, we strongly recommend it. To serve its purpose, the successive paragraphs below refresh some required aspects of mathematical language as used on the applied level.

1.1 Algebra

REAL NUMBERS

We typeset the set of:

- ▷ natural numbers (unsigned integers) as \mathbb{N} including zero,
- ▷ integer numbers as \mathbb{Z} including zero,
- ▷ rational numbers as \mathbb{Q} including zero,
- ▷ real numbers (floats) as \mathbb{R} including zero.

All the above make a chain of subsets: $\mathbb{N} \subset \mathbb{Z} \subset \mathbb{Q} \subset \mathbb{R}$.

To avoid possible confusion, we outline a brief glossary of mathematical terms. We recall that using the correct mathematical terms reflects correct mathematical thinking. Putting down ideas in the correct words is of major importance for profound insight.

Sets

- ▷ We recall writing all **subsets** in between braces, e.g. the **empty set** appears as $\{\}$.
- ▷ We define a **singleton** as any subset containing only one element, e.g. $\{5\} \subset \mathbb{N}$, as a subset of natural numbers.
- ▷ We define a **pair** as any subset containing just two elements, e.g. $\{115, -4\} \subset \mathbb{Z}$, as a subset of integers. In programming the boolean values *true* and *false* make up a pair $\{true, false\}$ called the boolean set which we typeset as \mathbb{B} .
- ▷ We define $\mathbb{Z}^- = \{\dots, -3, -2, -1\}$ whenever we need negative integers only. We express symbolically that -1234 is an **element** of \mathbb{Z}^- by typesetting $-1234 \in \mathbb{Z}^-$.
- ▷ We typeset the **set minus** operator to delete elements from a set by using a backslash, e.g. $\mathbb{N} \setminus \{0\}$ reading all natural numbers except zero, $\mathbb{Q} \setminus \mathbb{Z}$ meaning all pure rational numbers after all integer values left out and $\mathbb{R} \setminus \{0, 1\}$ expressing all real numbers apart from zero and one.

Calculation basics

operation	expression	a	b	c
to add	$a + b = c$	term	term	sum
to subtract	$a - b = c$	term	term	difference
to multiply	$a \cdot b = c$	factor	factor	product
to divide	$\frac{a}{b} = c, b \neq 0$	numerator	divisor or denominator	quotient or fraction
to exponentiate	$a^b = c$	base	exponent	power
to take the root	$\sqrt[b]{a} = c$	radicand	index	radical
return factorial	$n! = c$			n factorial

We define the **factorial** of a natural argument as the returned product of this argument multiplied with all natural numbers from this number n down to 1. Put in symbols:

$$n! = n \cdot (n-1) \cdot (n-2) \cdot \dots \cdot 3 \cdot 2 \cdot 1 \quad \text{restricted to } n \in \mathbb{N}$$

Furthermore we define $1! = 1$ and as well $0! = 1$.

Examples:

$$2! = 2 \cdot 1 = 2, \quad 3! = 3 \cdot 2 \cdot 1 = 6, \quad 4! = 4 \cdot 3 \cdot 2 \cdot 1 = 24.$$

We write the **opposite** of a real number r as $-r$, defined by the sum $r + (-r) = 0$. We typeset the **reciprocal** of a nonzero real number r as $\frac{1}{r}$ or r^{-1} , defined by the product $r \cdot r^{-1} = 1$.

We define **subtraction** as equivalent to adding the opposite: $a - b = a + (-b)$. We define **division** as equivalent to multiplying with the reciprocal: $a : b = a \cdot b^{-1}$.

When we mix operations we need to apply priority rules for them. There is a fixed priority list 'PEMDAS' in performing mixed operations in \mathbb{R} that can easily be memorised by 'Please Excuse My Dear Aunt Sally'.

- ▷ First process all that is delimited in between Parentheses,
- ▷ then Exponentiate,
- ▷ then Multiply and Divide from left to right,
- ▷ finally Add and Subtract from left to right.

Now we discuss the **distributive law** ruling within \mathbb{R} , which we define as threading a ‘superior’ operation over an ‘inferior’ operation. In conclusion, distributing requires two *different* operations.

Hence we distribute *exponentiating* over *multiplication* as in $(a \cdot b)^3 = a^3 \cdot b^3$. Likewise rules *multiplying* over *addition* as in $3 \cdot (a + b) = 3 \cdot a + 3 \cdot b$.



However we should never stumble on this ‘Staircase of Distributivity’ by going too fast:

$$(a + b)^3 \neq a^3 + b^3,$$

$$\sqrt{a + b} \neq \sqrt{a} + \sqrt{b},$$

$$\sqrt{x^2 + y^2} \neq x + y.$$

Fractions

A **fraction** is what we call any rational number written as $\frac{t}{n}$ given $t, n \in \mathbb{Z}$ and $n \neq 0$, wherein t is called the **numerator** and n the **denominator**. We define the reciprocal of a nonzero fraction $\frac{t}{n}$ as $\frac{1}{\frac{t}{n}} = \frac{n}{t}$ or as the power $\left(\frac{t}{n}\right)^{-1}$. We define the opposite fraction as $-\frac{t}{n} = \frac{-t}{n} = \frac{t}{-n}$. We summarise fractional arithmetic:

sum	$\frac{t}{n} + \frac{a}{b} = \frac{t \cdot b + n \cdot a}{n \cdot b}$
difference	$\frac{t}{n} - \frac{a}{b} = \frac{t \cdot b - n \cdot a}{n \cdot b}$
product	$\frac{t}{n} \cdot \frac{a}{b} = \frac{t \cdot a}{n \cdot b}$
division	$\frac{\frac{t}{a}}{\frac{n}{b}} = \frac{t}{n} \cdot \frac{b}{a}$
exponentiation	$\left(\frac{t}{n}\right)^m = \frac{t^m}{n^m}$
singular fractions	$\frac{1}{0} = \pm\infty$ infinity (see page 76)
	$\frac{0}{0} = ?$ indeterminate

Powers

We define a **power** as any real number written as g^m , wherein g is called its **base** and m its **exponent**. The opposite of g^m is simply $-g^m$. The reciprocal of g^m is $\frac{1}{g^m} = g^{-m}$, given $g \neq 0$.

According to the exponent type we distinguish between:

$$\begin{aligned} g^3 &= g \cdot g \cdot g & 3 \in \mathbb{N}, \\ g^{-3} &= \frac{1}{g^3} = \frac{1}{g \cdot g \cdot g} & -3 \in \mathbb{Z}, \\ g^{\frac{1}{3}} &= \sqrt[3]{g} = w \Leftrightarrow w^3 = g & \frac{1}{3} \in \mathbb{Q}, \\ g^0 &= 1 & g \neq 0. \end{aligned}$$

Whilst calculating powers we may have to:

$$\begin{aligned} \text{multiply} \quad & g^3 \cdot g^2 = g^{3+2} = g^5, \\ \text{divide} \quad & \frac{g^3}{g^2} = g^3 \cdot g^{-2} = g^{3-2} = g^1, \\ \text{exponentiate} \quad & (g^3)^2 = g^{3 \cdot 2} = g^6 \text{ them.} \end{aligned}$$

We insist on avoiding typesetting radicals like $\sqrt[7]{g^3}$ and strongly recommend their contemporary notation using radicand g and exponent $\frac{3}{7}$, consequently exponentiating g to $g^{\frac{3}{7}}$. We recall the fact that all square roots are non-negative numbers, $\sqrt{a} = a^{\frac{1}{2}} \in \mathbb{R}^+$ for $a \in \mathbb{R}^+$.

As well as knowing the above exponent types, understanding the above rules to calculate them is necessary for using powers successfully. We advise memorising the integer squares running from $1^2 = 1$, $2^2 = 4$, ..., up to $15^2 = 225$, $16^2 = 256$ and the integer cubes running from $1^3 = 1$, $2^3 = 8$, ..., up to $7^3 = 343$, $8^3 = 512$ in order to easily recognise them.

Recall that the only way out of any power is exponentiating with its reciprocal exponent. For this purpose we need to exponentiate both left hand side and right hand side of any given relation (see also paragraph 1.2).

Example: Find x when $\sqrt[7]{x^3} = 5$ by exponentiating this power.

$$x^{\frac{3}{7}} = 5 \Leftrightarrow \left(x^{\frac{3}{7}}\right)^{\frac{7}{3}} = (5)^{\frac{7}{3}} \Leftrightarrow x \approx 42.7494.$$

We emphasise the above strategy as the only successful one to free base x from its exponent, yielding its correct expression numerically approximated if we wish to.

Example: Find x when $x^2 = 5$ by exponentiating this power.


$$x^2 = 5 \Leftrightarrow (x^2)^{\frac{1}{2}} = (5)^{\frac{1}{2}} \text{ or } -(5)^{\frac{1}{2}} \Leftrightarrow x \approx 2.23607 \text{ or } -2.23607.$$

We recall the above double solution whenever we free base x from an *even* exponent, yielding their correct expression as accurately as we wish to.

Mathematical expressions

Composed mathematical expressions can often seem intimidating or cause confusion. To gain transparency in them, we firstly recall indexed variables which we define as subscripted to count them: $x_1, x_2, x_3, x_4, \dots, x_{99999}, x_{100000}, \dots$, and $\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4, \dots$. It is common practice in industrial research to use thousands of variables, so just picking unindexed characters would be insufficient. Taking our own alphabet as an example, it would only provide us with 26 characters.

We define finite expressions as composed of (mathematical) operations on objects (numbers, variables or structures). We can for instance analyse the expression $(3a+x)^4$ by drawing its **tree form**. This example reveals a Power having exponent 4 and a subexpression in its base. The base itself yields a sum of the variable x Plus another subexpression. This final subexpression shows the product 3 Times a .



Let us also evaluate this expression $(3a+x)^4$. Say $a = 1$, then we see our expression partly collapse to $(3+x)^4$. If, on top of this, we assign $x = 2$, our expression then finally turns to the numerical value $(3+2)^4 = 5^4 = 625$.

When we expand this power to its **pure sum expression** $81a^4 + 108a^3x + 54a^2x^2 + 12ax^3 + x^4$, we did nothing but *reshape* its **pure product expression** $(3a+x)^4$.

We warn that trying to solve this expression – which is not a relation – is completely in vain. Recall that inequalities, equations and systems of equations or inequalities are the only objects in the universe we can (try to) solve mathematically.

Relational operators

We also refresh the use of correct terms for inequalities and equations.

We define an **inequality** as any *variable* expression comparing a left hand side to a right hand side by applying the ‘is-(strictly)-less-than’ or by applying the ‘is-(strictly)-greater-than’ operator. For example, we can read $(3a+x)^4 \leq (b+4)(x+3)$ containing variables a, x, b . Consequently we may solve such inequality for any of the unknown quantities a, x or b .

We define an **equation** as any *variable* expression comparing a left hand side to a right hand side by applying the ‘is-equal-to’ operator. For example $(3a+x)^4 = (b+4)(x+3)$ is an equation containing variables a, x, b . Consequently we also may solve equations for any of the unknown quantities a, x or b .